

DYNAL

Encl. 1

PRODUCT LIST 1989



BEST AVAILABLE COPY

DYNABEADS™ products are based on extremely uniform, superparamagnetic polystyrene beads. Consisting of a maghemite (Fe_2O_3) containing core covered with a polymer, they have a smooth surface that is easily coated with antibodies or other selecting molecules. Combined with a magnet, Dynabeads make a unique tool in positive or negative separations.

Fields of applications include:
 Immunology, Tissue Typing, Cancer research,
 Transplantation medicine, Microbiology, Virology,
 DNA Technology and Clinical chemistry.

DYNABEADS UNCOATED

A. Immunomagnetic beads for cell separations. Uniform, superparamagnetic polystyrene beads with diameter 4.5 micron (c.v. < 5%).
 4×10^8 DYNABEADS per ml (30 mg per ml) in aqueous solution.

DYNABEADS M-450 Uncoated
 For physical adsorption of primary antibodies of the IgM class, or for customer's own secondary antibodies. Primary monoclonal antibodies of the IgG class should be bound to Dynabeads M-450 via a secondary antibody for optimal function.

DYNABEADS M-450 Tosylactivated
 For convenient chemical coupling of proteins or secondary antibodies of customers own choice.

B. Immunomagnetic beads for use in microbiology and immunoassays. Uniform superparamagnetic polystyrene beads with a polymer surface having only primary OH groups and with a diameter of 2.8 micron (c.v. < 3%).
 $6-7 \times 10^8$ DYNABEADS per ml (10 mg per ml) in aqueous solution.

NEW DYNABEADS M-280 Tosylactivated
 For convenient chemical coupling of proteins, peptides or secondary antibodies of customers own choice. Dynabeads M-280 are activated by use of p-toluene sulphonyl chloride and ready for coating through a simple incubation.

	Prod.no.	Volum:
	140.01	2
	140.02	10
	140.03	10 m
	140.04	
	142.03	
	142.04	

Enc. 4

BEST AVAILABLE COPY

NEW WORLDWIDE

006
NG.315
DYNAL R/S → D0441712060766
09:29 16/11/98

Promega Protocols and Applications Guide

Second Edition

Interdisciplinary
biat biology in
tools of
his effort, and is
; research
; molecular and cell

the growing
logy products.
at international
ions with
oviet Union.
; the global
; hard to develop
; consume. In 1989,
; program for our loan
; and recyclable
; boxes. To



© Copyright 1991 (March)
Promega Corporation
All rights reserved. Printed in U.S.A.

Part Number 1981

The Promega Protocols and Applications Guide
text pages are printed on recycled paper.

Promega Corporation
2810 Woods Hollow Road
Madison, WI 53711-5359 USA
Toll Free 800-356-9526
Telephone 608-274-4330
FAX 608-273-6967
Telex 25037092

800-356-9526


Promega

1.1.2. **Location, Purification and Labeling**

Nucleic Acid Detoxin, Future Research

הוּא־בָּהִר

Contents

سید ۳

Isolation of Total RNA

Section, Publishing, and Purification	147
X. 3'-End Labeling to Fill Accesses 3' Ends of Double-Stranded DNA	148
1. 3'-End Labeling with Terminal Transferase	148
A. Addition of [α-32P]dNTP "Tails" to 3' Termini of Single-Stranded DNA Primers	149
B. Addition of [α-32P]Cordycepin-5'-Triphosphate to 3' Termini of Single-Stranded DNA Primers	149
C. Determination of Percent Incorporation and Specific Activity	150
D. Gel Analysis	150
XI. 5'-End Labeling using T4 Polynucleotidic Kinase	151
A. Dephosphorylation Reaction	151
B. Kinase Reaction	151
C. Determination of Percent Incorporation	151
XII. References	151
XIII. Additional Nucleic Acid Labeling Literature Available from Promega	152
1. Immunoaffinity Isolation of β-Galactosidase Fusion Proteins using ProtaSorb[®] Fc⁺ Adherent Immunoaffinity Isolation of DNA Binding Proteins with the GRAB System	2
2. Immunoaffinity Isolation of Lambda Expression Libraries with the ProtoBlot[™] Immunoaffinity Screening System	2
3. Activity Purification of DNA Binding Proteins with the GRAB System	3

I. Magnetic Particle Separation of Molecules

The attachment of nucleic acids to solid support has found many applications in the field of molecular biology. One common application of immobilized nucleic acids is oligo(dT) cellulose purification of messenger RNA (mRNA) by hybridization to the 3' polyadenine tail (4). Recent years, however, have witnessed the emergence of paramagnetic particles as the solid support of choice for many affinity purification protocols. Paramagnetic particles which have iron oxide into submicron sized particles which have no magnetic field but form a magnetic dipole when exposed to a magnetic field. The use of paramagnetic particles eliminates the need for traditional column chromatography, centrifugation, or any other special equipment. These particles have been successfully used in the development of immunoassays (5), diagnostic assays (6), and for measuring RNA in cell lysates using cDNA-tailed capture probes (7).

Promega has extended the use of paramagnetic particles to the affinity purification of polyadenylated mRNA with the PolyATract™ system and to cDNA synthesis and cloning with the Capture Clone™ system. Unlike procedures which use direct coupling of probes to paramagnetic particles (6,7), these systems use a biotinylated digonucleotide probe to hybridize in solution to the targeted nucleic acid. The hybrids are then captured using covalently coupled streptavidin paramagnetic particles. This approach combines the speed and efficiency of solution hybridization with the convenience and speed (<1 minute) of magnetic separation.

used for purified blots, cDNA syn-

specific oligonucleotide probe used. For *Drosophila* *d17*, the calculated binding capacity is roughly 1 nM probe captured/mg SA-PMPs.